

## TIMP3

**Reactivity:** Human Mouse Rat

**Tested applications:** WB IHC IF

**Recommended Dilution:** WB 1:500 - 1:2000 IHC 1:50 - 1:200 IF 1:50 - 1:200

**Calculated MW:** 24kDa

**Observed MW:** Refer to Figures

**Immunogen:**

Recombinant protein of human TIMP3

**Storage Buffer:**

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

**Concentration:**

b

**Synonym:**

HSMRK222; K222; K222TA2; SFD;TIMP3;

**Catalog #:** A1511

**Antibody Type:**

Polyclonal Antibody

**Species:** Rabbit

**Gene ID:** 7078

**Isotype:** IgG

**Swiss Prot:** P35625

**Purity:** Affinity purification

For research use only.

**Background:**

TIMPs are members of the family of tissue inhibitor of matrix metalloproteinases (MMPs) that includes TIMP1, TIMP2, TIMP3, and TIMP4. The main function of TIMPs is their inhibitory effect on MMPs. TIMPs irreversibly inactivate MMPs by direct binding to their catalytic zinc cofactor and resultant inhibition of proteinase function (1,2). In addition to MMP inhibition, TIMPs have also been shown to interact with various membrane receptors on the cell surface. Some of these interactions include: TIMP1 with CD63, TIMP2 with 31 integrin, and TIMP3 with VEGFR2, all of which result in distinct cellular effects (3). TIMPs are involved in a wide variety of biological functions, such as tumor angiogenesis and progression (4,5), wound healing, and vascular remodeling (6,7). Mutations in TIMP3 are associated with Sorsby's fundus dystrophy (8,9).

**To place an order, please [Click HERE](#).**