www.neobiolab.com info@neobiolab.com 888.754.5670, +1 617.500.7103 United States 0800.088.5164, +44 020.8123.1558 United Kingdom

## HDAC2

Reactivity: Human Mouse Rat

Tested applications: WB IHC IF

Recommended Dilution:WB 1:500 - 1:2000 IHC 1:50 - 1:200 IF 1:20 - 1:50

Calculated MW:55kDa

Observed MW:Refer to Figures

Immunogen:

A synthetic peptide of human HDAC2

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol,

pH7.3.

Concentration:

fip

Synonym:

RPD3; YAF1;

## Background:

Acetylation of the histone tail causes chromatin to adopt an "open" conformation, allowing increased accessibility of transcription factors to DNA. The identification of histone acetyltransferases (HATs) and their large multiprotein complexes has yielded important insights into how these enzymes regulate transcription (1,2). HAT complexes interact with sequence-specific activator proteins to target specific genes. In addition to histones, HATs can acetylate nonhistone proteins, suggesting multiple roles for these enzymes (3). In contrast, histone deacetylation promotes a "closed" chromatin conformation and typically leads to repression of gene activity (4). Mammalian histone deacetylases can be divided into three classes on the basis of their similarity to various yeast deacetylases (5). Class I proteins (HDACs 1, 2, 3, and 8) are related to the yeast Rpd3-like proteins, those in class II (HDACs 4, 5, 6, 7, 9, and 10) are related to yeast Hda1-like proteins, and class III proteins are related to the yeast protein Sir2. Inhibitors of HDAC activity are now being explored as potential therapeutic cancer agents (6,7).HDAC1 and HDAC2 are highly homologous and are involved in histone deacetylation, chromatin remodeling and transcriptional repression (8-10). Both proteins are found together in numerous complexes including the nucleosome remodeling and deacetylation complex (NuRD), MeCP1, and the mSin3A corepressor complex.

To place an order, please Click HERE.



Catalog #:A2084

**Antibody Type:** 

Polyclonal Antibody

Species: Rabbit

Gene ID:3066 Isotype:IgG

Swiss Prot:Q92769

Purity: Affinity purification

For research use only.





