www.neobiolab.com info@neobiolab.com 888.754.5670, +1 617.500.7103 United States 0800.088.5164, +44 020.8123.1558 United Kingdom

IL 10 Human, His

Description: Interleukin-10 Human Recombinant produced in E.Coli is single, a non-glycosylated, Polypeptide chain containing 181 amino acids fragment (19-178) and having a total molecular mass of 20.94kDa with a 20 amino acids N-Terminal His tag. The IL-10 His-Tag protein is purified by proprietary chromatographic techniques.

Catalog #:CYPS-493

For research use only.

Synonyms:B-TCGF, CSIF, TGIF, IL-10, IL10A, MGC126450, MGC126451, Cytokine synthesis inhibitory factor.

Source: Escherichia Coli.

Physical Appearance: Sterile Filtered clear solution.

Amino Acid Sequence: MGSSHHHHHH SSGLVPRGSH MSPGQGTQSE NSCTHFPGNL PNMLRDLRDA FSRVKTFFQM KDQLDNLLLK ESLLEDFKGY LGCQALSEMI QFYLEEVMPQ AENQDPDIKA HVNSLGENLK TLRLRRRCH RFLPCENKSK AVEQVKNAFN KLQEKGIYKA MSEFDIFINY IEAYMTMKIR N.

Purity: Greater than 95.0% by SDS-PAGE.

Formulation:

Interleukin-10 His is supplied in 20mM Tris-HCl pH-8 and 20% glycerol.

Stability:

Store at 4°C if entire vial will be used within 2-4 weeks. Store, frozen at -20°C for longer periods of time. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Please avoid freeze thaw cycles.

Usage:

NeoBiolab's products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

Introduction:

Interleukin-10 is a multifunctional cytokine produced by a variety of cell types including T helper cells activated B cells and activated macrophages. Interleukin-10 regulates immune-mediated inflammation and modulates the function of cells such as lymphocytes, monocytes, natural killer cells and dendritic cells.

To place an order, please Click HERE.





