

BD 2 Human

Description: Beta Defensin-2 Human Recombinant produced in E.Coli is a single, non-glycosylated polypeptide chain containing 41 amino acids and having a molecular mass of 4.3 kDa. The BD-2 is purified by proprietary chromatographic techniques.

Synonyms: BD-2, hBD-2, Defensin beta 2, Skin-antimicrobial peptide 1, SAP1, DEFB4, DEFB102, DEFB2, DEFB4P, Beta-defensin 2.

Source: Escherichia Coli.

Physical Appearance: Sterile Filtered White lyophilized (freeze-dried) powder.

Amino Acid Sequence: GIGDPVTCLK SGAICHPVFC PRRYKQIGTC GLPGTKCCKK P.

Purity: Greater than 98.0% as determined by: (a) Analysis by RP-HPLC. (b) Analysis by SDS-PAGE.

Formulation:

The Human BD-2 was lyophilized from a 0.2

Stability:

Lyophilized Beta Defensin-2 Recombinant although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution BD-2 should be stored at 4°C between 2-7 days and for future use below -18°C. Please prevent freeze-thaw cycles.

Usage:

NeoBiolab's products are furnished for LABORATORY RESEARCH USE ONLY. They may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

Solubility:

It is recommended to reconstitute the lyophilized Beta Defensin-2 in sterile 18M-cm H₂O not less than 100µg/ml, which can then be further diluted to other aqueous solutions.

Introduction:

The Defensin family are highly similar in their protein sequence and are microbicidal & cytotoxic peptides made by neutrophils. Beta Defensin-1 is an antimicrobial peptide having the resistance of epithelial surfaces to microbial colonization. Beta Defensin-1 has close proximity to Defensin Alpha-1 and has been implicated in the pathogenesis of cystic fibrosis. Skin of patients having atopic dermatitis patients and mycosis fungoides (non-lesional and lesional) show lower human Beta Defensin-1 mRNA expression and higher human Beta Defensin-2 and human Beta Defensin-3 mRNA expression. BBeta Defensin is highly expressed by epithelial cells. Beta-defensin 1 may play a role in the pathogenesis of severe sepsis. Variation in human Beta Defensin-1 contributes to asthma diagnosis, with apparent gender-specific effects. Human Beta Defensin-3 is a dimer, while Human BD-1 and Human BD-2 are monomeric. The expression of Human BD1 is correlated with induction profiles in gingival keratinocytes. The level of expression of human DEFB1 mRNA is lower than that of human BD3 and human BD-2 in reconstructed epidermis. Human BD1 is down-regulated in human prostatic and renal carcinomas.

Biological Activity:


www.neobiolab.com

info@neobiolab.com

888.754.5670, +1 617.500.7103 United States

0800.088.5164, +44 020.8123.1558 United Kingdom

Determined by the ability to chemoattract human dendritic immature cells at a concentration of 10-100ng/ml.



To place an order, please [Click HERE](#).

Catalog #:CYP5-578

For research use only.

© 2015 NeoScientific



neobiolab | [Term](#) | [Quote/Order](#) | [Services](#) | [Products](#) | [Support](#) | [Corporate](#) | [Contact Us](#)

Copyrights @ 2013 NEO Group 245 First Street, 18th Floor, Cambridge MA 02142 888.754.5670 (toll-free)



Request a
CATALOG
[CLICK HERE](#)