

HGF B Human

Description: HGF-B Human Recombinant produced in E.Coli is a single, non-glycosylated, Polypeptide chain containing 234 amino acids fragment (495-728) having a molecular weight of 34kDa and fused with a 4.5kDa amino-terminal hexahistidine tag. The HGF-B is purified by proprietary chromatographic techniques.

Catalog #: CYPs-672

For research use only.

Synonyms: Scatter Factor, SF, Hepatopoietin, HPTA, HGF, HGFB, F-TCF, DFNB39, Hepatocyte growth factor, Hepatocyte growth factor beta chain.

Source: Escherichia Coli.

Physical Appearance: Sterile Filtered clear solution.

Amino Acid Sequence:

VVNGIPTRTNIGWMVSLRYRNKHICGGSLIKESWVLTARQCPSRDLKDYEAWLGIHVDVHGRGDE
KCKQVLNVSQLVYGPEGSDDLVMKLARPAVLDDFVSTIDLPNYGCTIPEKTSVYGWGYTGTLIN
YDGLLRVAHLYIMGNEKCSQHHRGKVTLNESEICAGAEKIGSGPCEGDYGGPLVCEQHKMRMVL
GVIVPGRGCAIPNRPGIFVRVAYYAKWIKILTYKVPQS.

Purity: Greater than 95.0% as determined by SDS-PAGE.

Formulation:

HGF-B protein is supplied in 10mM TrisHCl (pH.8), 50% glycerol

Stability:

Store at 4°C if entire vial will be used within 2-4 weeks. Store, frozen at -20°C for longer periods of time. Please avoid freeze thaw cycles.

Usage:

NeoBiolab's products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

Introduction:

Hepatocyte Growth Factor (HGF) is a multifunctional growth factor which regulates both cell growth and cell motility. It exerts a strong mitogenic effect on hepatocytes and primary epithelial cells. HGF synergizes with Interleukin-3 and GM-CSF to stimulate colony formation of hematopoietic progenitor cells in vitro and may, therefore, also modulate hematopoiesis. HGF is secreted as a single inactive polypeptide which is cleaved by serine proteases into a 69kDa Alpha chain and 34kDa Beta chain. A disulfide bond linking the alpha and beta chains produces the active heterodimeric molecule.

To place an order, please [Click HERE](#).