

## Trypsin Porcine

**Description:** Recombinant Porcine Trypsin is free from any animal and human sources. Recombinant Porcine Trypsin expressed in Yeast and purified by standard chromatography techniques. Recombinant Porcine Trypsin is free from foreign enzymes such as carboxypeptidase A & chymotrypsin. Recombinant Human Trypsin is free from protease inhibitors such as PMSF and EDTA.

**Catalog #:** PRPS-794

For research use only.

**Source:** Pichia Pastoris.

**Physical Appearance:** Sterile Filtered clear liquid solution.

**Amino Acid Sequence:**

VGGYTCAANSIPYQVSLNSGSHFCGGSLNSQWVVSAAHCYKSRIQVRLGEHNIDVLEGNEQFIN  
AAKIITHPNFNGNTLDNDIMLIKSSPATLNSRVATVSLPRSCAAAGTECLISGWGNTKSSGSSYPS  
LLQCLKAPVLSDSCKSSYPGQITGNMICVGFLEGGKDSCQGDGGPVVCNGQLQGIVSWGYG  
CAQKNKPGVYTKVCNYYVNWIIQQTIAAN.

**Formulation:**

The Porcine Trypsin (2.98mg/ml) is formulated with 1mM HCl and 20mM CaCl<sub>2</sub>, pH-3.

**Stability:**

Recombinant Porcine Trypsin should be stored at 2-8°C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA).

**Usage:**

NeoBiolab's products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

**Introduction:**

Trypsin (EC3.4.21.4) is part of the serine protease family. Trypsin cleaves lysine and arginine at the C-terminal side of the peptide. The hydrolysis rate is slower if an acidic residue is on either sides of the cleavage site and no cleavage occurs if a proline residue is on the carboxyl side of the cleavage site. Trypsin optimum pH is pH-7 to 9. Trypsin will also hydrolyze ester and amide linkages of synthetic derivatives of amino acids such as: benzoyl L-arginine ethyl ester (BAEE), p-toluenesulfonyl- L-arginine methyl ester (TAME), tosyl-L-arginine methyl ester, N-benzoyl-L-arginine p-nitroanilide (BAPNA), L-lysyl-p-nitroanilide, and benzoyl-L-tyrosine ethyl ester (BTEE). Serine protease inhibitors that inhibit recombinant trypsin include TLCK (N-p-tosyl-L-lysine chloromethyl ketone), PMSF (phenylmethanesulfonyl fluoride), benzamidine, soybean trypsin inhibitor, and ovomucoid.

**Biological Activity:**

3,394 BAEE units/mg powder.

**To place an order, please [Click HERE](#).**